

## Structure-Based Discovery of A<sub>2A</sub> Adenosine Receptor Ligands

Jens Carlsson,<sup>†</sup> Lena Yoo,<sup>‡</sup> Zhan-Guo Gao,<sup>‡</sup> John J. Irwin,<sup>†</sup> Brian K. Shoichet,<sup>\*,†</sup> and Kenneth A. Jacobson<sup>\*,‡</sup>

<sup>†</sup>Department of Pharmaceutical Chemistry, University of California, 1700 4th Street, Box 2550, San Francisco, California 94158, and

<sup>‡</sup>Molecular Recognition Section, Laboratory of Bioorganic Chemistry, National Institute of Diabetes and Digestive and Kidney Diseases, National Institutes of Health, Bethesda, Maryland 20892

Received February 23, 2010

The recent determination of X-ray structures of pharmacologically relevant GPCRs has made these targets accessible to structure-based ligand discovery. Here we explore whether novel chemotypes may be discovered for the A<sub>2A</sub> adenosine receptor, based on complementarity to its recently determined structure. The A<sub>2A</sub> adenosine receptor signals in the periphery and the CNS, with agonists explored as anti-inflammatory drugs and antagonists explored for neurodegenerative diseases. We used molecular docking to screen a 1.4 million compound database against the X-ray structure computationally and tested 20 high-ranking, previously unknown molecules experimentally. Of these 35% showed substantial activity with affinities between 200 nM and 9 μM. For the most potent of these new inhibitors, over 50-fold specificity was observed for the A<sub>2A</sub> versus the related A<sub>1</sub> and A<sub>3</sub> subtypes. These high hit rates and affinities at least partly reflect the bias of commercial libraries toward GPCR-like chemotypes, an issue that we attempt to investigate quantitatively. Despite this bias, many of the most potent new ligands were novel, dissimilar from known ligands, providing new lead structures for modulation of this medically important target.

### Introduction

G-protein-coupled receptors (GPCRs<sup>a</sup>) are a large family of transmembrane proteins that signal intracellularly after binding an extracellular ligand. These receptors share a similar topology, with seven transmembrane helices, but recognize a wide range of different signaling molecules. GPCRs have been intensely studied as pharmaceutical targets, and over 40% of marketed drugs act through them.<sup>1</sup> Until recently, a missing link to deeper understanding of GPCRs has been a lack of atomic resolution structural information. With the recent advent of several X-ray crystal structures of pharmacologically relevant GPCRs<sup>2–5</sup> it has for the first time become possible to leverage high-resolution structures for ligand discovery against these targets.<sup>6</sup>


Among the new GPCR structures is that of the A<sub>2A</sub> adenosine receptor (AR).<sup>5</sup> There are four subtypes of the AR (A<sub>1</sub>, A<sub>2A</sub>, A<sub>2B</sub>, and A<sub>3</sub>), and they are activated by extracellular adenosine in response to organ stress or tissue damage. The A<sub>2A</sub> AR signals in both the periphery and the CNS, with agonists explored as anti-inflammatory drugs and antagonists explored for neurodegenerative diseases, e.g., Parkinson's disease.<sup>7–11</sup> Although access to high resolution structural data is a crucial step toward atomic-level understanding of GPCRs,

the lack of structures has certainly not been an obstacle for successful ligand discovery. For several decades, classical ligand-based medicinal chemistry approaches have been used to identify thousands of AR ligands. Almost all known AR agonists are derivatives of the cognate ligand (**1–3**, Chart 1), whereas antagonists are more diverse. Two large classes of AR antagonists are xanthines, with members such as caffeine (**4**) and theophylline (**5**), and adenine derivatives such as **6** (ZM241385<sup>12</sup>), which is bound to the A<sub>2A</sub> AR binding site in the crystallographic structure (Chart 1, Figure 1A). Despite considerable medicinal chemistry efforts and the wide range of possible therapeutic applications for AR ligands, there are only a few approved drugs targeting this receptor.<sup>8,11</sup> Consequently, there remains an ongoing need for new subtype selective agonists and antagonists of this target.

Here, we wished to investigate whether we could find new A<sub>2A</sub> AR ligand chemotypes by using structure-based molecular docking to screen a large and putatively unbiased library of small molecules, looking for those that complement the receptor structure. Docking evaluates the complementarity of small molecules to a receptor binding site of known structure<sup>13–18</sup> and can in principle discover new chemotypes, dissimilar to previous ligands, that nevertheless fit the binding site well. Such chemotypes might provide new routes for modulation of this key target. Methodologically, we wanted to explore what the hit rate of a structure-based (docking) screen against the A<sub>2A</sub> AR might be. In docking screens against the β<sub>2</sub> adrenergic GPCR, a hit rate of 24% had been observed.<sup>19–23</sup> A docking “hit” is a molecule that binds to the target at a relevant concentration, and a docking “hit rate” is the number of compounds that bind divided by the number of compounds experimentally tested. For the β<sub>2</sub> adrenergic


\*To whom correspondence should be addressed. For B.K.S.: phone, 415-514-4126; fax, 415-514-4260; e-mail, shoichet@cgl.ucsf.edu. For K.A.J.: phone, 301-496-9024; fax, 301-480-8422; e-mail, kajacobs@helix.nih.gov.

<sup>a</sup> Abbreviations: GPCR, G-protein-coupled receptor; AR, adenosine receptor; CNS, central nervous system; SEA, similarity ensemble approach; PDB, Protein Data Bank; WOMBAT, World of Molecular Bioactivity; CHO, Chinese hamster ovary; DMEM, Dulbecco's modified Eagle medium; PKA, protein kinase A; DLS, dynamic light scattering.



**Figure 1.** Binding mode of the cocrystallized ligand **6** (A) and the predicted binding modes of the seven ligands discovered in the docking screen (B–H). The  $A_{2A}$  AR binding site is shown in white ribbons with the side chains of Glu169 and Asn253 in sticks. In (A) the cocrystallized ligand **6** is shown using orange carbon atoms. In (B–H), the crystallographic ligand is shown using blue lines and the docking poses for the ligands are depicted with orange carbon atoms. Black dotted lines indicate hydrogen bonds. The compounds are (B) **7**, (C) **8**, (D) **9**, (E) **10**, (F) **11**, (G) **12**, and (H) **13**.

**Chart 1.** Structures of Known Agonists (**1–3**) and Antagonists (**4–6**) of the  $A_{2A}$  Adenosine Receptor



receptor, where the affinity of the best docking hit was 9 nM, both were unusually high. We wished to understand whether this would be true for this second GPCR and why this might be so. To investigate this, we docked a library of 1.4 million small molecules to the crystal structure of the  $A_{2A}$  AR. From the top-scoring molecules, 20 were selected on the basis of their fit to the binding site and chemical diversity. Here, we present the experimental evaluation of these molecules and assess why GPCRs appear to be particularly suitable targets for structure-based ligand discovery.

## Methods

**Preparation of the Molecular Docking Screen.** All docking calculations were carried out with the program DOCK 3.5.54<sup>16,17,24</sup> using a 2.6 Å crystallographic structure of the  $A_{2A}$  adenosine receptor in complex with an antagonist (**6**)<sup>12</sup> (PDB accession code 3EML<sup>5</sup>). The receptor structure was prepared by removing all non-protein atoms and the intracellular T4-lysozyme insertion. The protonation states of ionizable residues were set to the most probable in aqueous solution at pH 7.

The flexible-ligand sampling algorithm in DOCK3.5.54 superimposes atoms of the docked molecule onto binding site matching spheres, which indicate putative ligand atom positions.<sup>16,17</sup> In the case of the  $A_{2A}$  AR, 45 matching spheres were used, and these were either based on the atoms of the crystallographic ligand or positioned manually. The spheres were also labeled for chemical matching based on the local receptor environment.<sup>25</sup> The degree of ligand sampling is determined by the bin size, bin size overlap, and distance tolerance. These three parameters were set to 0.4, 0.3, and 1.5 Å, respectively, for both the binding site matching spheres and the docked molecules. For ligand conformations passing an initial steric filter, a physics-based scoring function is used to evaluate the fit to the receptor binding site. For the best scoring conformation of each docked molecule, 100 steps of rigid-body minimization are carried out. The score for each conformation is calculated as the sum of the receptor–ligand electrostatic and van der Waals interaction energy, corrected for ligand desolvation. These three terms are evaluated from precalculated grids. The three-dimensional map of the electrostatic potential in the binding site was prepared using the program Delphi.<sup>26</sup> In this calculation, partial charges from the united atom AMBER force field<sup>27</sup> were used

for all receptor atoms except the side chain amide of Asn253, for which the dipole moment was increased to favor hydrogen bonding to this residue (we have adopted this technique of increasing local dipoles on a few polar residues in the active site without changing their formal charges extensively in past studies).<sup>28,29</sup> The program CHEMGRID was used to generate a van der Waals grid, which is based on a united atom version of the AMBER force field.<sup>30</sup> The desolvation penalty for a ligand conformation is estimated from a precalculated transfer free energy of the molecule between solvents of dielectrics 78 and 2. The desolvation energy is obtained by weighting the transfer free energy with a scaling factor that reflects the degree of burial of the ligand in the receptor binding site.<sup>31,32</sup>

The ZINC leadlike set was prepared by filtering a large library of commercially available compounds using the criteria  $\log P < 3.5$ , molecular weight of  $< 350$ , and number of rotatable bonds of  $\leq 7$ .<sup>33</sup> Each molecule has been prepared for docking by pregenerating up to 1000 conformations using the program OMEGA.<sup>34</sup> Partial atomic charges and transfer free energies have been calculated using AMSOL,<sup>35,36</sup> and van der Waals parameters have been derived from an all-atom AMBER potential.<sup>37</sup>

**Similarity and Library Bias Calculations.** Similarity calculations were carried out with the program Pipeline Pilot<sup>38</sup> using the Tanimoto coefficient and ECFP4 fingerprints. For each of the docking-discovered ligands, the Tanimoto similarity to all annotated A<sub>1</sub>, A<sub>2A</sub>, A<sub>2B</sub>, and A<sub>3</sub> AR ligands with  $K_i \leq 10 \mu\text{M}$  in the World of Molecular Bioactivity (WOMBAT 2006.2)<sup>39</sup> and ChEMBL (a StARlite 2009 prerelease version)<sup>40</sup> databases was calculated. The number of molecules in the ZINC leadlike database that are similar to known ligands of the ARs, adrenergic receptors, adenylyl cyclases, and AmpC  $\beta$ -lactamase was predicted with the similarity ensemble approach (SEA) using ECFP4 fingerprints.<sup>41</sup> Ligands ( $K_i \leq 10 \mu\text{M}$ ) that are annotated to ARs, adrenergic receptors, and adenylyl cyclases were extracted from the WOMBAT database. Ligands for AmpC  $\beta$ -lactamase were extracted from refs 28, 29, and 42. For each ligand set, the number of leadlike molecules in ZINC that have a SEA  $P$  value better than  $10^{-10}$  was calculated. The predicted compounds were then postfiltered for molecules that match the molecular weight and formal charge ranges of the known ligands.

**A<sub>2A</sub> AR Receptor Binding and Functional Assay.** Binding assays at three hAR subtypes were carried out using standard radioligands<sup>43–45</sup> and membrane preparations from Chinese hamster ovary (CHO) cells (A<sub>1</sub> and A<sub>3</sub>) or human embryonic kidney (HEK293) cells (A<sub>2A</sub>) stably expressing a hAR subtype.<sup>46,47</sup> A functional assay at the A<sub>2A</sub>AR consisted of stimulation of cAMP production<sup>48,49</sup> in A<sub>2A</sub>AR-expressing HEK293 cells. [<sup>3</sup>H]R-N<sup>6</sup>-(2-phenylisopropyl)adenosine ([<sup>3</sup>H]R-PIA, 42.6 Ci/mmol) was obtained from Moravek Biochemicals (Brea, CA). [<sup>3</sup>H](2-[*p*-(2-Carboxyethyl)phenylethylamino]-5'-*N*-ethylcarboxamidoadenosine) ([<sup>3</sup>H]CGS21680, 40.5 Ci/mmol) and [<sup>125</sup>I]N<sup>6</sup>-(4-amino-3-iodobenzyl)adenosine-5'-*N*-methyluronamide ([<sup>125</sup>I]I-AB-MECA, 2200 Ci/mmol) were purchased from Perkin-Elmer Life and Analytical Science (Boston, MA). Test compounds were prepared as 5 mM stock solutions in DMSO and stored frozen at  $-20^\circ\text{C}$ .

**Cell Culture and Membrane Preparation.** CHO cells stably expressing the recombinant hA<sub>1</sub> and hA<sub>3</sub>Rs, and HEK-293 cells stably expressing the hA<sub>2A</sub>AR were cultured in Dulbecco's modified Eagle medium (DMEM) and F12 (1:1) supplemented with 10% fetal bovine serum, 100 units/mL penicillin, 100  $\mu\text{g}/\text{mL}$  streptomycin, and 2  $\mu\text{mol}/\text{mL}$  glutamine. In addition, 800  $\mu\text{g}/\text{mL}$  Geneticin was added to the A<sub>2A</sub> media, while 500  $\mu\text{g}/\text{mL}$  hygromycin was added to the A<sub>1</sub> and A<sub>3</sub> media. After being harvested, cells were homogenized and suspended in PBS. Cells were then centrifuged at 240g for 5 min, and the pellet was resuspended in 50 mM Tris-HCl buffer (pH 7.5) containing 10 mM MgCl<sub>2</sub>. The suspension was homogenized and was then















ultracentrifuged at 14330g for 30 min at  $4^\circ\text{C}$ . The resultant pellets were resuspended in Tris buffer and incubated with adenosine deaminase (3 units/mL) for 30 min at  $37^\circ\text{C}$ . The suspension was homogenized with an electric homogenizer for 10 s, pipetted into 1 mL vials, and then stored at  $-80^\circ\text{C}$  until the binding experiments were conducted. The protein concentration was measured using the BCA protein assay kit from Thermo Scientific Pierce Protein Research Products (Rockford, IL).<sup>50</sup>

**Binding Assays.** The tested compounds were purchased from six different vendors (Enamine, ChemDiv, ChemBridge, Vitas-M, Pharmeks, and Asinex). The vendors had verified that each compound had  $\geq 95\%$  purity by liquid chromatography–mass spectrometry (LC–MS) or nuclear magnetic resonance (NMR) experiments. Into each tube in the binding assay was added 50  $\mu\text{L}$  of increasing concentrations of the test ligand in Tris-HCl buffer (50 mM, pH 7.5) containing 10 mM MgCl<sub>2</sub>, 50  $\mu\text{L}$  of the appropriate agonist radioligand, and finally 100  $\mu\text{L}$  of membrane suspension. For the A<sub>1</sub>AR (22  $\mu\text{g}$  of protein/tube) the radioligand used was [<sup>3</sup>H]R-PIA (final concentration of 3.5 nM). For the A<sub>2A</sub>AR (20  $\mu\text{g}$ /tube) the radioligand used was [<sup>3</sup>H]CGS21680 (10 nM). For the A<sub>3</sub>AR (21  $\mu\text{g}$ /tube) the radioligand used was [<sup>125</sup>I]I-AB-MECA (0.34 nM). Nonspecific binding was determined using a final concentration of 10  $\mu\text{M}$  unlabeled 5'-*N*-ethylcarboxamidoadenosine (NECA, **2**) diluted with the buffer. The mixtures were incubated at  $25^\circ\text{C}$  for 60 min in a shaking water bath. Binding reactions were terminated by filtration through Brandel GF/B filters under a reduced pressure using a M-24 cell harvester (Brandel, Gaithersburg, MD). Filters were washed three times with 3 mL of 50 mM ice-cold Tris-HCl buffer (pH 7.5). Filters for A<sub>1</sub> and A<sub>2A</sub>AR binding were placed in scintillation vials containing 5 mL of Hydrofluor scintillation buffer and counted using a Perkin Elmer liquid scintillation analyzer (Tri-Carb 2810TR). Filters for A<sub>3</sub>AR binding were counted using a Packard Cobra II  $\gamma$ -counter. The  $K_i$  values were determined using GraphPad Prism for all assays.

**Cyclic AMP Accumulation Assay.** Intracellular cyclic AMP (cAMP) levels were measured with a competitive protein binding method.<sup>48,49</sup> CHO293 cells that expressed the recombinant human A<sub>2A</sub>AR were harvested by trypsinization. After centrifugation and resuspension in medium, cells were planted in 24-well plates in 1.0 mL of medium. After 24 h, the medium was removed and cells were washed three times with 1 mL of DMEM containing 50 mM HEPES, pH 7.4. Cells were then treated with the test compound in the presence of rolipram (10  $\mu\text{M}$ ) and adenosine deaminase (3 units/mL), and incubation was continued for an additional 1 h. The reaction was terminated by removing the supernatant, and cells were lysed upon the addition of 200  $\mu\text{L}$  of 0.1 M ice-cold HCl. The cell lysate was resuspended and stored at  $-20^\circ\text{C}$ . For determination of cyclic AMP production, protein kinase A (PKA) was incubated with [<sup>3</sup>H]cyclic AMP (2 nM) in K<sub>2</sub>HPO<sub>4</sub>/EDTA buffer (K<sub>2</sub>HPO<sub>4</sub>, 150 mM; EDTA, 10 mM), 20  $\mu\text{L}$  of the cell lysate, and 30  $\mu\text{L}$  of 0.1 M HCl or 50  $\mu\text{L}$  of cyclic AMP solution (0–16 pmol/200  $\mu\text{L}$  for standard curve). Bound radioactivity was separated by rapid filtration through Whatman GF/C filters and washed once with cold buffer. Bound radioactivity was measured by liquid scintillation spectrometry.

**Counterscreen for Colloidal Inhibition.** To control for artifactual inhibition by colloidal aggregation, we looked for particle formation by Dynamic Light Scattering (DLS) and by inhibition of two counterscreen enzymes, cruzain and AmpC  $\beta$ -lactamase. Concentrated DMSO stocks of compounds were diluted with filtered 50 mM KPi, pH 7.0. Measurements were made using a DynaPro MS/X (Wyatt Technology) with a 55 mW laser at 826.6 nm. The laser power was 100%, and the detector angle was  $90^\circ$ . Cruzain assays were performed in 100 mM sodium acetate, pH 5.5, containing 5 mM DTT with and without 0.1% Triton X-100. Compounds were incubated with 0.8 nM cruzain for 5 min, and reactions were initiated by

**Table 1.** Ligand Structures and Experimental Data for the Seven Hits from the Docking Screen against the A<sub>1</sub>, A<sub>2A</sub>, and A<sub>3</sub> ARs

ID (Rank) <sup>a</sup>	Ligand structure	K <sub>i</sub> (μM) or % displacement at 20 μM <sup>b</sup>			Most similar known adenosine receptor ligand <sup>c</sup>	T <sub>c</sub> <sup>d</sup>
		A <sub>1</sub>	A <sub>2A</sub>	A <sub>3</sub>		
7 (8)		15%	8.8 ± 1.5	8%		0.52
8 (83)		4%	3.6 ± 2.4	37%		0.37
9 (88)		19%	0.2 ± 0.0	3.0 ± 0.2		0.37
10 (114)		5%	2.7 ± 1.0	28%		0.32
11 (198)		25%	0.2 ± 0.1	18%		0.30
12 (483)		6%	1.2 ± 0.2	17%		0.61
13 (500)		12%	1.3 ± 0.7	15%		0.31

<sup>a</sup> Rank in the docking screen. <sup>b</sup> Measured in three independent experiments. <sup>c</sup> The most similar compound annotated to ARs in WOMBAT<sup>39</sup> and ChEMBL.<sup>40</sup> <sup>d</sup> Tanimoto similarity coefficient to the closest annotated adenosine receptor ligand from ECFP4 fingerprints.

adding the fluorogenic substrate Z-Phe-Arg-aminomethylcoumarin (Z-FR-AMC). The final reaction volume was 200 μL with cruzain at 0.4 nM and ZF-R-AMC at 2.5 μM. Final DMSO concentrations were 0.5%. To measure enzyme inhibition, the increase in fluorescence (excitation wavelength of 355 nm, emission wavelength of 460 nm) was recorded for 5 min in a microtiter plate spectrofluorimeter (Molecular Devices, Flex-Station). Assays were performed in duplicate in 96-well plates, with controls measuring enzyme activity in the presence of DMSO. Activity was measured for seven different concentrations for each compound. Inhibition of AmpC β-lactamase was measured for the two best compounds identified here, compounds **9** and **11**, to complement the cruzain assay results. Assays were performed in 50 mM potassium phosphate, pH 7.0. Compounds were incubated with 1 nM β-lactamase for 5 min, and reactions were initiated by adding the substrate CENTA to a final concentration 92 μM. The final reaction volume was 1 mL. To measure enzyme inhibition, the increase in absorbance at 405 nm was recorded for 5 min in a UV-vis spectrophotometer (Agilent). Assays were performed in duplicate in 1 mL cuvettes, with controls measuring enzyme activity in the presence of DMSO. Activity was measured at 10 μM, 0.1% DMSO.

## Results and Discussion

**Molecular Docking Screen and Compound Selection.** The program DOCK3.5.54<sup>16,17</sup> was used to screen 1.4 million commercially available “leadlike” molecules from the ZINC<sup>33</sup>

database against the orthosteric site of the A<sub>2A</sub> AR. On average, each molecule was sampled in 5000 orientations and, for each that fit, an average of 16 500 conformations; the receptor was held rigid. Each molecule was scored for electrostatic and van der Waals complementarity corrected for ligand desolvation. Molecules typically overlapped with the geometry of the crystallized antagonist, making a mixture of polar and hydrophobic interactions, packing deeply into the site (Figure 1). The 500 top-ranking molecules (Table S1, Supporting Information), 0.035% of the docking prioritized library, were analyzed visually for features that are not taken into account in the docking calculation. This is a standard procedure for all our docking screens in which each of the molecules is inspected for novelty, physical properties, and binding energy contributions that are not included in the docking scoring function. For example, compound **9** (Table 1) was chosen both because it complemented the site well and because there were several molecules with the same scaffold in the top 500 list of compounds (e.g., the molecules ranked 7, 31, 59, 104, 115, 135, 137, 172, 184, 199, and 278 in Table S1, Supporting Information). From this evaluation, 20 compounds representing diverse molecules (within the context of the top 500-ranking molecules) were prioritized for experimental testing.

**Radioligand Displacement Assays and Docking Hit Rate.** The 20 compounds selected from the docking screen were tested for binding in a radioligand displacement assay. Seven



**Table 3.** Target Library Bias and Docking Hit Rates

target	no. of ZINC molecules similar to known ligands <sup>a</sup>	representative DOCK screen	
		hit rate <sup>b</sup> (%)	best potency <sup>c</sup> (nM)
adenosine receptors	4240	35	200
adrenergic receptors	4146	24	9 <sup>19</sup>
adenylyl cyclases	565	4	50000 <sup>55</sup>
AmpC $\beta$ -lactamase	545	2–5	26000 <sup>29</sup>


<sup>a</sup>ZINC leadlike molecules with at least  $10^{-10}$  *P* values to annotated target ligands in WOMBAT using the similarity ensemble approach (SEA).  
<sup>b</sup>(Number of true ligands)/(number of predictions tested experimentally).  
<sup>c</sup>The affinity of the ligand with the best potency from the docking screen.

control calculations. Asn253 is conserved in all four AR receptor subtypes and has also been found to be a crucial interaction partner for both agonists and antagonists in mutagenesis studies.<sup>52</sup>

Whereas all the seven ligands are previously uncharacterized for the A<sub>2A</sub> AR, some of them bear known chemotypes. To quantify their novelty, or lack of it, we calculated the similarity of each molecule to 7500 known AR ligands from the WOMBAT and ChEMBL databases using pairwise Tanimoto coefficients (*T<sub>c</sub>*, ECFP4 fingerprints) (Table 1). Ligand **12** resembles members of the xanthine class of antagonists, while compound **7** resembles certain quinazoline<sup>53</sup> ligands. Conversely, whereas compounds **10**, **11**, and **13** do conserve several moieties with known ligands, they also differ substantially from them, with *T<sub>c</sub>* values of 0.3 to the closest annotated ligand. Nevertheless, they complement the site well both sterically and electrostatically (Figure 1). The potency of these molecules suggests that they may merit further study as new lead families for antagonists of the A<sub>2A</sub> AR.

To determine the efficacy of the compounds, their ability to inhibit intracellular cAMP production induced by agonist **3** was tested. No stimulation of cAMP production was detected for any of the molecules, while a clear displacement of agonist function was observed for the two most potent compounds, **9** and **11** (Figure 3). All seven of the new ligands are thus almost certainly A<sub>2A</sub> antagonists, as is the cocrystallized ligand. Intriguingly, this efficacy bias was also observed in the docking screens against the  $\beta_2$  adrenergic receptor, where only inverse agonists were found against the structure crystallized with the inverse agonist carazolol. In docking to rigid GPCR structures, the protein conformation may bias the screen toward molecules with the same efficacy as the cocrystallized ligand. This represents a challenge to our ability to exploit these structures for mechanisms of action, such as agonism, not represented in the experimental structure. To further investigate this, we determined the rank of two agonists, **1** and **2** (Chart 1), that were present in the set of commercially available molecules screened against the A<sub>2A</sub> AR binding site. Whereas multiple known antagonists would have been ranked among the top 500 molecules in the docking screen, these two agonists were ranked 951 057 and 919 993, respectively.

**Is There Library Bias toward GPCR Chemotypes in Chemical Libraries?** Returning to one of the questions that motivated this study, the structure-based screen against the A<sub>2A</sub> AR returned a diverse set of ligands dissimilar to those previously characterized, as well as several similar to known ligands, and did so with a hit rate of 35%. Not only is this hit rate much higher than we have come to expect for enzyme targets screened with the same approach, but the new antagonists were also close to 100-fold more potent than we have come to expect for our docking “hits”. Furthermore, these results are strikingly similar to those observed in docking



**Figure 3.** Functional assay based on measuring the production of cAMP for **3** (control), a potent A<sub>2A</sub> AR agonist, with or without 10  $\mu$ M **9** or **11**. The dose–response curve is shifted for both compounds, as expected in the case of competitive antagonistic inhibition. The % activation refers to production of cAMP normalized to the effect of **3** at 100  $\mu$ M.

screens against the  $\beta_2$  adrenergic GPCR.<sup>19,20</sup> To what may these unusually high hit rates and affinities be attributed?

Family A GPCRs like the A<sub>2A</sub> AR are the targets for a substantial fraction of marketed drugs, and this partly reflects the quality of their sites for specific recognition of small molecules. Largely buried from bulk solvent, these sites can almost completely enclose a “druglike” molecule and can do so with a mixture of nonpolar and polar interactions. Consequently, a large and sustained medicinal chemistry effort has focused on these targets, and by now even putatively unbiased libraries, like ZINC, have become populated with molecules bearing “GPCR-like” chemotypes. This also reflects a bias toward naturally occurring molecules in our screening libraries.<sup>54</sup> Indeed, Kolb et al. estimated that there were 3–12 times as many small molecules that were similar to GPCR ligands in the ZINC leadlike set compared to other common drug targets such as kinases, proteases, and ligand-gated ion channels.<sup>19</sup> To make this comparison more specific and relevant to the adenosine receptor versus other docking targets that we ourselves have worked, we investigated the library bias in ZINC for the adrenergic and adenosine receptors together with two other targets for which we have observed much lower hit rates and affinities<sup>19,29,55</sup> (Table 3). The number of molecules in the ZINC leadlike set that are similar to the ligands of these targets was estimated using SEA,<sup>54</sup> insisting on a *P* value of  $10^{-10}$  or better; acceptable molecules had also to resemble the annotated ligands in their physical properties (see Methods). Over 4000 small molecules resemble ligands annotated to the ARs and adrenergic receptors in the WOMBAT<sup>39</sup> database, almost 10-fold more than found for the enzymes adenylyl cyclase and AmpC  $\beta$ -lactamase, against which, correspondingly, our docking hit rates and affinities have been 10- to 100-fold lower. Thus, it is the convolution of the high “ligand-ability” of the orthosteric sites and the many GPCR-like chemotypes in

our libraries that makes the adrenergic and adenosine receptors so fruitful for structure-based techniques.

**Note Added after Initial Review of This Paper.** After this paper was submitted for review, a paper by Abagyan, Stevens, and colleagues appeared that also targeted the A<sub>2A</sub> AR for novel inhibitor discovery, also using a molecular docking screen.<sup>56</sup> As here, Abagyan et al. also observed a very substantial hit rate with high affinities (indeed, both the affinities and hit rates were slightly better than those we observe). Whereas the compounds discovered in the two screens were substantially different, the observation of the high hit rates and, by screening standards, high affinity “hits” is consistent with the high “ligand-ability” of the class A GPCRs and the fortuitous library bias toward them, and this is among the important conclusions of this study.

**Acknowledgment.** This work is supported by NIH Grant GM59957 (to B.K.S.), NIDDK Intramural Research Program (to K.A.J.), and a fellowship from the Knut and Alice Wallenberg Foundation (to J.C.). We thank A. Doak for aggregation assays and members of the Shoichet lab for docking “hit-list” evaluation. We thank Tudor Oprea for access to the WOMBAT database and John Overington for a prerelease version of the ChEMBL database.

**Supporting Information Available:** Table S1 of structures of the 500 top-ranking molecules from the docking screen. This material is available free of charge via the Internet at <http://pubs.acs.org>.

## References

- Overington, J. P.; Al-Lazikani, B.; Hopkins, A. L. How many drug targets are there? *Nat. Rev. Drug Discovery* **2006**, *5*, 993–996.
- Cherezov, V.; Rosenbaum, D. M.; Hanson, M. A.; Rasmussen, S. G. F.; Thian, F. S.; Kobilka, T. S.; Choi, H. J.; Kuhn, P.; Weis, W. I.; Kobilka, B. K.; Stevens, R. C. High-resolution crystal structure of an engineered human beta(2)-adrenergic G protein-coupled receptor. *Science* **2007**, *318*, 1258–1265.
- Rosenbaum, D. M.; Cherezov, V.; Hanson, M. A.; Rasmussen, S. G. F.; Thian, F. S.; Kobilka, T. S.; Choi, H. J.; Yao, X. J.; Weis, W. I.; Stevens, R. C.; Kobilka, B. K. GPCR engineering yields high-resolution structural insights into beta(2)-adrenergic receptor function. *Science* **2007**, *318*, 1266–1273.
- Warne, T.; Serrano-Vega, M. J.; Baker, J. G.; Moukhametzianov, R.; Edwards, P. C.; Henderson, R.; Leslie, A. G. W.; Tate, C. G.; Schertler, G. F. X. Structure of a beta(1)-adrenergic G-protein-coupled receptor. *Nature* **2008**, *454*, 486–491.
- Jaakola, V. P.; Griffith, M. T.; Hanson, M. A.; Cherezov, V.; Chien, E. Y. T.; Lane, J. R.; IJzerman, A. P.; Stevens, R. C. The 2.6 angstrom crystal structure of a human A(2A) adenosine receptor bound to an antagonist. *Science* **2008**, *322*, 1211–1217.
- Congreve, M.; Marshall, F. The impact of GPCR structures on pharmacology and structure-based drug design. *Br. J. Pharmacol.* **2010**, *159*, 986–996.
- Moro, S.; Gao, Z. G.; Jacobson, K. A.; Spalluto, G. Progress in the pursuit of therapeutic adenosine receptor antagonists. *Med. Res. Rev.* **2006**, *26*, 131–159.
- Jacobson, K. A.; Gao, Z. G. Adenosine receptors as therapeutic targets. *Nat. Rev. Drug Discovery* **2006**, *5*, 247–264.
- Sebastiao, A. M.; Ribeiro, J. A. Adenosine receptors and the central nervous system. *Handb. Exp. Pharmacol.* **2009**, 471–534.
- Blackburn, M. R.; Vance, C. O.; Morschl, E.; Wilson, C. N. Adenosine receptors and inflammation. *Handb. Exp. Pharmacol.* **2009**, 215–269.
- Cristalli, G.; Muller, C. E.; Volpini, R. Recent developments in adenosine A<sub>2A</sub> receptor ligands. *Handb. Exp. Pharmacol.* **2009**, 59–98.
- Poucher, S. M.; Keddie, J. R.; Singh, P.; Stogdall, S. M.; Caulkett, P. W. R.; Jones, G.; Collis, M. G. The in-vitro pharmacology of Zm-241385, a potent, nonxanthine, a(2a) selective adenosine receptor antagonist. *Br. J. Pharmacol.* **1995**, *115*, 1096–1102.
- Degen, J.; Rarey, M. FlexNovo: structure-based searching in large fragment spaces. *ChemMedChem* **2006**, *1*, 854–868.
- Jones, G.; Willett, P.; Glen, R. C.; Leach, A. R.; Taylor, R. Development and validation of a genetic algorithm for flexible docking. *J. Mol. Biol.* **1997**, *267*, 727–748.
- Kairys, V.; Fernandes, M. X.; Gilson, M. K. Screening drug-like compounds by docking to homology models: a systematic study. *J. Chem. Inf. Model.* **2006**, *46*, 365–379.
- Lorber, D. M.; Shoichet, B. K. Flexible ligand docking using conformational ensembles. *Protein Sci.* **1998**, *7*, 938–950.
- Lorber, D. M.; Shoichet, B. K. Hierarchical docking of databases of multiple ligand conformations. *Curr. Top. Med. Chem.* **2005**, *5*, 739–749.
- Zavodszky, M. I.; Kuhn, L. A. Side-chain flexibility in protein–ligand binding: the minimal rotation hypothesis. *Protein Sci.* **2005**, *14*, 1104–1114.
- Kolb, P.; Rosenbaum, D. M.; Irwin, J. J.; Fung, J. J.; Kobilka, B. K.; Shoichet, B. K. Structure-based discovery of beta(2)-adrenergic receptor ligands. *Proc. Natl. Acad. Sci. U.S.A.* **2009**, *106*, 6843–6848.
- Sabio, M.; Jones, K.; Topiol, S. Use of the X-ray structure of the beta(2)-adrenergic receptor for drug discovery. Part 2: Identification of active compounds. *Bioorg. Med. Chem. Lett.* **2008**, *18*, 5391–5395.
- de Graaf, C.; Rognan, D. Selective structure-based virtual screening for full and partial agonists of the beta 2 adrenergic receptor. *J. Med. Chem.* **2008**, *51*, 4978–4985.
- Katritch, V.; Reynolds, K. A.; Cherezov, V.; Hanson, M. A.; Roth, C. B.; Yeager, M.; Abagyan, R. Analysis of full and partial agonists binding to beta(2)-adrenergic receptor suggests a role of transmembrane helix V in agonist-specific conformational changes. *J. Mol. Recognit.* **2009**, *22*, 307–318.
- Reynolds, K. A.; Katritch, V.; Abagyan, R. Identifying conformational changes of the beta(2) adrenoceptor that enable accurate prediction of ligand/receptor interactions and screening for GPCR modulators. *J. Comput.-Aided Mol. Des.* **2009**, *23*, 273–288.
- Kuntz, I. D.; Blaney, J. M.; Oatley, S. J.; Langridge, R.; Ferrin, T. E. A geometric approach to macromolecule–ligand interactions. *J. Mol. Biol.* **1982**, *161*, 269–288.
- Shoichet, B. K.; Kuntz, I. D. Matching chemistry and shape in molecular docking. *Protein Eng.* **1993**, *6*, 723–732.
- Nicholls, A.; Honig, B. A rapid finite-difference algorithm, utilizing successive over-relaxation to solve the Poisson–Boltzmann equation. *J. Comput. Chem.* **1991**, *12*, 435–445.
- Weiner, S. J.; Kollman, P. A.; Case, D. A.; Singh, U. C.; Ghio, C.; Alagona, G.; Profeta, S.; Weiner, P. A new force-field for molecular mechanical simulation of nucleic-acids and proteins. *J. Am. Chem. Soc.* **1984**, *106*, 765–784.
- Babaoglu, K.; Simeonov, A.; Lrwin, J. J.; Nelson, M. E.; Feng, B.; Thomas, C. J.; Cancian, L.; Costi, M. P.; Maltby, D. A.; Jadhav, A.; Inglese, J.; Austin, C. P.; Shoichet, B. K. Comprehensive mechanistic analysis of hits from high-throughput and docking screens against beta-lactamase. *J. Med. Chem.* **2008**, *51*, 2502–2511.
- Powers, R. A.; Morandi, F.; Shoichet, B. K. Structure-based discovery of a novel, noncovalent inhibitor of AmpC beta-lactamase. *Structure* **2002**, *10*, 1013–1023.
- Meng, E. C.; Shoichet, B. K.; Kuntz, I. D. Automated docking with grid-based energy evaluation. *J. Comput. Chem.* **1992**, *13*, 505–524.
- Shoichet, B. K.; Leach, A. R.; Kuntz, I. D. Ligand solvation in molecular docking. *Proteins: Struct., Funct., Genet.* **1999**, *34*, 4–16.
- Wei, B. Q. Q.; Baase, W. A.; Weaver, L. H.; Matthews, B. W.; Shoichet, B. K. A model binding site for testing scoring functions in molecular docking. *J. Mol. Biol.* **2002**, *322*, 339–355.
- Irwin, J. J.; Shoichet, B. K. ZINC—a free database of commercially available compounds for virtual screening. *J. Chem. Inf. Model.* **2005**, *45*, 177–182.
- Bostrom, J.; Greenwood, J. R.; Gottfries, J. Assessing the performance of OMEGA with respect to retrieving bioactive conformations. *J. Mol. Graphics Modell.* **2003**, *21*, 449–462.
- Chambers, C. C.; Hawkins, G. D.; Cramer, C. J.; Truhlar, D. G. Model for aqueous solvation based on class IV atomic charges and first solvation shell effects. *J. Phys. Chem.* **1996**, *100*, 16385–16398.
- Li, J. B.; Zhu, T. H.; Cramer, C. J.; Truhlar, D. G. New class IV charge model for extracting accurate partial charges from wave functions. *J. Phys. Chem. A* **1998**, *102*, 1820–1831.
- Weiner, S. J.; Kollman, P. A.; Nguyen, D. T.; Case, D. A. An all atom force-field for simulations of proteins and nucleic-acids. *J. Comput. Chem.* **1986**, *7*, 230–252.
- <http://accelrys.com/products/scitegic/>.
- Olah, M.; Mracec, M.; Ostopovic, L.; Rad, R.; Bora, A.; Hadaruga, N.; Olah, I.; Banda, M.; Simon, Z.; Mracec, M.; Oprea, T. I. WOMBAT: World of Molecular Bioactivity. In *Cheminformatics in Drug Discovery*; Oprea, T. I., Ed.; Wiley-VCH: Weinheim, Germany, 2005; pp 221–239.
- <http://www.ebi.ac.uk/chembl>.

- (41) Keiser, M. J.; Roth, B. L.; Armbruster, B. N.; Ernsberger, P.; Irwin, J. J.; Shoichet, B. K. Relating protein pharmacology by ligand chemistry. *Nat. Biotechnol.* **2007**, *25*, 197–206.
- (42) Tondi, D.; Morandi, F.; Bonnet, R.; Costi, M. P.; Shoichet, B. K. Structure-based optimization of a non-beta-lactam lead results in inhibitors that do not up-regulate beta-lactamase expression in cell culture. *J. Am. Chem. Soc.* **2005**, *127*, 4632–4639.
- (43) Jarvis, M. F.; Schulz, R.; Hutchison, A. J.; Do, U. H.; Sills, M. A.; Williams, M. [H-3] Cgs-21680, a selective A2 adenosine receptor agonist directly labels A2-receptors in rat-brain. *J. Pharmacol. Exp. Ther.* **1989**, *251*, 888–893.
- (44) Klotz, K. N.; Lohse, M. J.; Schwabe, U.; Cristalli, G.; Vittori, S.; Grifantini, M. 2-Chloro-*N*-6-[H-3]cyclopentyladenosine ([H-3]Ccpa), a high-affinity agonist radioligand for A1 adenosine receptors. *Naunyn-Schmiedeberg's Arch. Pharmacol.* **1989**, *340*, 679–683.
- (45) Olah, M. E.; Gallorodriguez, C.; Jacobson, K. A.; Stiles, G. L. I-125 4-aminobenzyl-5'-*N*-methylcarboxamidoadenosine, a high-affinity radioligand for the rat a(3) adenosine receptor. *Mol. Pharmacol.* **1994**, *45*, 978–982.
- (46) Englert, M.; Qwitterer, U.; Klotz, K. N. Effector coupling of stably transfected human A(3) adenosine receptors in CHO cells. *Biochem. Pharmacol.* **2002**, *64*, 61–65.
- (47) Jacobson, K. A.; Park, K. S.; Jiang, J. L.; Kim, Y. C.; Olah, M. E.; Stiles, G. L.; Ji, X. D. Pharmacological characterization of novel A(3) adenosine receptor-selective antagonists. *Neuropharmacology* **1997**, *36*, 1157–1165.
- (48) Nordstedt, C.; Fredholm, B. B. A modification of a protein-binding method for rapid quantification of camp in cell-culture supernatants and body-fluid. *Anal. Biochem.* **1990**, *189*, 231–234.
- (49) Post, S. R.; Ostrom, R. S.; Insel, P. A. Biochemical methods for detection and measurement of cyclic AMP and adenylyl cyclase activity. *Methods Mol. Biol.* **2000**, *126*, 363–374.
- (50) Bradford, M. M. Rapid and sensitive method for quantitation of microgram quantities of protein utilizing principle of protein–dye binding. *Anal. Biochem.* **1976**, *72*, 248–254.
- (51) McGovern, S. L.; Helfand, B. T.; Feng, B.; Shoichet, B. K. A specific mechanism of nonspecific inhibition. *J. Med. Chem.* **2003**, *46*, 4265–4272.
- (52) Kim, J. H.; Wess, J.; Vanrhee, A. M.; Schoneberg, T.; Jacobson, K. A. Site-directed mutagenesis identifies residues involved in ligand recognition in the human a(2a) adenosine receptor. *J. Biol. Chem.* **1995**, *270*, 13987–13997.
- (53) Webb, T. R.; Lvovskiy, D.; Kim, S. A.; Ji, X. D.; Melman, N.; Linden, J.; Jacobson, K. A. Quinazolines as adenosine receptor antagonists: SAR and selectivity for A(2B) receptors. *Bioorg. Med. Chem.* **2003**, *11*, 77–85.
- (54) Hert, J.; Irwin, J. J.; Laggner, C.; Keiser, M. J.; Shoichet, B. K. Quantifying biogenic bias in screening libraries. *Nat. Chem. Biol.* **2009**, *5*, 479–483.
- (55) Soelaiman, S.; Wei, B. Q.; Bergson, P.; Lee, Y. S.; Shen, Y.; Mrksich, M.; Shoichet, B. K.; Tang, W. J. Structure-based inhibitor discovery against adenylyl cyclase toxins from pathogenic bacteria that cause anthrax and whooping cough. *J. Biol. Chem.* **2003**, *278*, 25990–25997.
- (56) Katritch, V.; Jaakola, V. P.; Lane, J. R.; Lin, J.; Ijzerman, A. P.; Yeager, M.; Kufareva, I.; Stevens, R. C.; Abagyan, R. Structure-based discovery of novel chemotypes for adenosine A(2A) receptor antagonists. *J. Med. Chem.* **2010**, *53*, 1799–1809.